

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
26 May 2005 (26.05.2005)

PCT

(10) International Publication Number
WO 2005/047315 A2

(51) International Patent Classification⁷: C07K 14/025

(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(21) International Application Number:
PCT/US2004/037372

(22) International Filing Date:
10 November 2004 (10.11.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/519,211 12 November 2003 (12.11.2003) US

(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BRYAN, Janine, T. [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). BROWNLOW, Michelle, K. [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). SCHULTZ, Loren, D. [US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). WANG, Xin-Min [CN/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). JANSEN, Kathrin, U. [DE/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: OPTIMIZED EXPRESSION OF HPV 58 L1 IN YEAST

Transmission EM of VLPs Composed of HPV 58 L1 R Protein Molecules



(57) Abstract: Synthetic DNA molecules encoding the HPV58 L1 protein are provided. Specifically, the present invention provides polynucleotides encoding HPV58 L1 protein, wherein said polynucleotides are codon-optimized for high level expression in a yeast cell. The synthetic molecules may be used to produce HPV58 virus-like particles (VLPs), and to produce vaccines and pharmaceutical compositions comprising the HPV58 VLPs. The vaccines of the present invention provide effective immunoprophylaxis against papillomavirus infection through neutralizing antibody and cell-mediated immunity and are also useful for treatment of existing HPV infections.

WO 2005/047315 A2

21571
#8

TITLE OF THE INVENTION

OPTIMIZED EXPRESSION OF HPV 58 L1 IN YEAST

FIELD OF THE INVENTION

The present invention relates generally to the prevention and/or therapy of human papillomavirus (HPV) infection. More specifically, the present invention relates to synthetic polynucleotides encoding HPV58 L1 protein, and to recombinant vectors and hosts comprising said polynucleotides. This invention also relates to HPV58 virus-like particles (VLPs), wherein the VLPs are produced by expressing recombinant HPV 58 L1 or L1 + L2 in yeast cells and to their use in vaccines and pharmaceutical compositions for preventing and treating HPV infections.

BACKGROUND OF THE INVENTION

There are more than 80 types of human papillomavirus (HPV), many of which have been associated with a wide variety of biological phenotypes, from benign proliferative warts to malignant carcinomas (for review, *see* McMurray et al., *Int. J. Exp. Pathol.* 82(1): 15-33 (2001)). HPV6 and HPV11 are the types most commonly associated with benign warts, nonmalignant condyloma acuminata and/or low-grade dysplasia of the genital or respiratory mucosa. HPV16 and HPV18 are the high-risk types most frequently associated with *in situ* and invasive carcinomas of the cervix, vagina, vulva and anal canal. More than 90% of cervical carcinomas are associated with infections of HPV16, HPV18 or the less prevalent oncogenic types HPV31, -33, -45, -52 and -58 (Schiffman et al., *J. Natl. Cancer Inst.* 85(12): 958-64 (1993)). The observation that HPV DNA is detected in more than 90% of cervical cancers provides strong epidemiological evidence that HPVs cause cervical carcinoma.

Papillomaviruses are small (50-60 nm), nonenveloped, icosahedral DNA viruses that encode up to eight early and two late genes. The open reading frames (ORFs) of the viral genomes are designated E1 to E7, and L1 and L2, where "E" denotes early and "L" denotes late. L1 and L2 code for virus capsid proteins, while the E genes are associated with functions such as viral replication and cellular transformation.

The L1 protein is the major capsid protein and has a molecular weight of 55-60 kDa. The L2 protein is the minor capsid protein. Immunological data suggest that most of the L2 protein is internal to the L1 protein in the viral capsid. Both the L1 and L2 proteins are highly conserved among different papillomaviruses.

Expression of the L1 protein or a combination of the L1 and L2 proteins in yeast, insect cells, mammalian cells or bacteria leads to self-assembly of virus-like particles (VLPs) (for review, *see* Schiller and Roden, in *Papillomavirus Reviews: Current Research on Papillomaviruses*; Lacey, ed. Leeds, UK: Leeds Medical Information, pp 101-12 (1996)).

5 VLPs are morphologically similar to authentic virions and are capable of inducing high titres of neutralizing antibodies upon administration into animals or humans. Because VLPs do not contain the potentially oncogenic viral genome, they present a safe alternative to the use of live virus in HPV vaccine development (for review, *see* Schiller and Hidesheim, *J. Clin. Virol.* 19: 67-74 (2000)). For this reason, the L1 and L2 genes have been identified as

10 immunological targets for the development of prophylactic and therapeutic vaccines for HPV infection and disease.

HPV vaccine development and commercialization have been hindered by difficulties associated with obtaining high expression levels of capsid proteins in successfully transformed host organisms, limiting the production of purified protein. Therefore, despite

15 the identification of wild-type nucleotide sequences encoding HPV L1 proteins such as HPV58 L1 protein, it would be highly desirable to develop a readily renewable source of crude HPV L1 protein that utilizes HPV58 L1-encoding nucleotide sequences that are optimized for expression in the intended host cell. Additionally, it would be useful to produce large quantities of HPV58 L1 VLPs having the immunity-conferring properties of

20 the native proteins for use in vaccine development.

SUMMARY OF THE INVENTION

The present invention relates to compositions and methods to elicit or enhance immunity to the protein products expressed by HPV58 L1 genes. Specifically, the

25 present invention provides polynucleotides encoding HPV58 L1 protein, wherein the polynucleotides have been codon-optimized for high level expression in a yeast cell. The present invention further provides HPV58 virus-like particles (VLPs), wherein said VLPs are produced by expressing recombinant HPV58 L1 or L1 + L2 in yeast cells, and discloses use of HPV58 VLPs in pharmaceutical compositions and vaccines for the prevention and/or

30 treatment of HPV-associated cancer.

The present invention relates to synthetic DNA molecules encoding the HPV58 L1 protein. The codons of the synthetic molecules are designed so as to use the codons preferred by a yeast cell. The synthetic molecules may be used as a source of HPV58 L1 protein, which may self-assemble into VLPs. Said VLPs may be used in a VLP-based

35 vaccine.

An exemplary embodiment of the present invention comprises a synthetic nucleic acid molecule which encodes the HPV58 L1 protein as set forth in SEQ ID NO:2, said nucleic acid molecule comprising a sequence of nucleotides that is codon-optimized for high-level expression in a yeast cell. In preferred embodiments, the nucleic acid comprises a
5 sequence of nucleotides as set forth in SEQ ID NO:1 (designated herein "58 L1 R sequence").

Also provided are recombinant vectors and recombinant host cells, both prokaryotic and eukaryotic, which contain the nucleic acid molecules disclosed throughout this specification. In a preferred embodiment of the present invention, the host cell is a yeast
10 cell.

The present invention also relates to a process for expressing an HPV58 L1 protein in a recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid encoding an HPV58 L1 protein into a yeast host cell; and (b) culturing the yeast host cell under conditions which allow expression of said HPV58 L1 protein.

15 The present invention further relates to a process for expressing an HPV58 L1 protein in a recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid molecule encoding an HPV58 L1 protein into a yeast host cell; wherein the nucleic acid molecule is codon-optimized for optimal expression in the yeast host cell and; (b) culturing the yeast host cell under conditions which allow expression of said HPV58 L1
20 protein.

In preferred embodiments of this aspect of the invention, the nucleic acid comprises a sequence of nucleotides as set forth in SEQ ID NO:1.

This invention also relates to HPV58 virus-like particles (VLPs) which are produced in yeast cells, methods of producing HPV58 VLPs, and methods of using HPV58
25 VLPs.

In a preferred embodiment of the invention, the yeast is selected from the group consisting of: *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Pichia pastoris*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, and *Schizosaccharomyces pombe*.

Another aspect of this invention is an HPV58 VLP, wherein the VLP is
30 produced by recombinant expression of HPV58 L1 or HPV58 L1 + L2 in a yeast cell.

Yet another aspect of this invention is an HPV58 VLP which comprises an HPV58 L1 protein produced by a codon-optimized HPV58 L1 gene. In an exemplary embodiment of this aspect of the invention, the codon-optimized HPV58 L1 gene comprises a sequence of nucleotides as set forth in SEQ ID NO:1.

This invention also provides a method for inducing an immune response in an animal comprising administering HPV58 virus-like particles to the animal. In a preferred embodiment, the HPV58 VLPs are produced by a codon-optimized gene.

Yet another aspect of this invention is a method of preventing or treating HPV-associated cervical cancer comprising administering to a mammal a vaccine comprising HPV58 VLPs. In a preferred embodiment of this aspect of the invention, the HPV58 VLPs are produced in yeast.

This invention also relates to a vaccine comprising HPV58 virus-like particles (VLPs), wherein the HPV58 VLPs are produced in yeast.

In an alternative embodiment of this aspect of the invention, the vaccine further comprises VLPs of at least one additional HPV type. The at least one additional HPV type may be any HPV type of interest, including any HPV type described in the art or those subsequently identified. In a preferred embodiment, the HPV type is a type that is associated with a clinical phenotype such as warts or cervical cancer. In a further preferred embodiment, the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV45, HPV51, HPV52, HPV55, HPV56, HPV59, and HPV68.

This invention also relates to pharmaceutical compositions comprising HPV 58 virus-like particles and a pharmaceutically acceptable carrier, wherein the HPV58 VLPs are produced in yeast. Further, this invention relates to pharmaceutical compositions comprising HPV58 VLPs and VLPs of at least one additional HPV type. In a preferred embodiment, the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV45, HPV51, HPV52, HPV55, HPV56, HPV59, and HPV68.

As used throughout the specification and in the appended claims, the singular forms "a," "an," and "the" include the plural reference unless the context clearly dictates otherwise.

As used throughout the specification and appended claims, the following definitions and abbreviations apply:

The term "promoter" refers to a recognition site on a DNA strand to which the RNA polymerase binds. The promoter forms an initiation complex with RNA polymerase to initiate and drive transcriptional activity. The complex can be modified by activating sequences termed "enhancers" or "upstream activating sequences" or inhibiting sequences termed "silencers".

The term "vector" refers to some means by which DNA fragments can be introduced into a host organism or host tissue. There are various types of vectors including plasmids, viruses (including adenovirus), bacteriophages and cosmids.

The term "cassette" refers to a nucleotide or gene sequence that is to be expressed from a vector, for example, the nucleotide or gene sequence encoding the HPV 58 L1 protein. In general, a cassette comprises a gene sequence inserted into a vector which, in some embodiments, provides regulatory sequences for expressing the nucleotide or gene sequence. In other embodiments, the nucleotide or gene sequence provides the regulatory sequences for its expression. In further embodiments, the vector provides some regulatory sequences and the nucleotide or gene sequence provides other regulatory sequences. For example, the vector can provide a promoter for transcribing the nucleotide or gene sequence and the nucleotide or gene sequence provides a transcription termination sequence. The regulatory sequences which can be provided by the vector include, but are not limited to, enhancers, transcription termination sequences, splice acceptor and donor sequences, introns, ribosome binding sequences, and poly(A) addition sequences.

The designations "58 L1 wild-type sequence" and "58 L1 wt sequence" refer to the HPV58 L1 sequence disclosed herein as SEQ ID NO:3. Although the HPV 58 L1 wild-type sequence has been described previously, it is not uncommon to find minor sequence variations between DNAs obtained from clinical isolates. Therefore, a representative 58 L1 wild-type sequence was isolated from clinical samples previously shown to contain HPV 58 DNA (see EXAMPLE 1). The 58 L1 wild-type sequence was used as a reference sequence to compare the codon-optimized 58 L1 sequences disclosed herein (see FIGURE 1).

The designations "HPV 58 L1 R" and "58 L1 R" refer to an exemplary synthetic HPV58 L1 nucleotide sequence (SEQ ID NO:1), disclosed herein, wherein the sequence was rebuilt so that it comprises codons that are preferred for high-level expression by a yeast cell.

The term "effective amount" means sufficient vaccine composition is introduced to produce the adequate levels of the polypeptide, so that an immune response results. One skilled in the art recognizes that this level may vary.

A "conservative amino acid substitution" refers to the replacement of one amino acid residue by another, chemically similar, amino acid residue. Examples of such conservative substitutions are: substitution of one hydrophobic residue (isoleucine, leucine, valine, or methionine) for another; substitution of one polar residue for another polar residue of the same charge (e.g., arginine for lysine; glutamic acid for aspartic acid).

The term "mammalian" refers to any mammal, including a human being.

"VLP" or "VLPs" mean(s) virus-like particle or virus-like particles.

"Synthetic" means that the HPV58 L1 gene was created so that it contains a sequence of nucleotides that is not the same as the sequence of nucleotides present in the designated naturally occurring wild-type HPV58 L1 gene (58 L1 wt, SEQ ID NO:3). As stated above, synthetic molecules are provided herein comprising a sequence of nucleotides comprising codons that are preferred for expression by yeast cells. The synthetic molecules provided herein encode the same amino acid sequences as the wild-type HPV58 L1 gene (SEQ ID NO:2).

10

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 shows a sequence alignment comparing nucleotides that were altered in the synthetic HPV58 L1 gene of the present invention (SEQ ID NO:1, indicated as "58 L1 R") (See EXAMPLE 2). The reference sequence is the 58 L1 wild-type sequence (SEQ ID NO:3, indicated as "58 L1 wt"; see EXAMPLE 1). Altered nucleotides are indicated at their corresponding location. Nucleotide number is contained within the parentheses. Identical nucleotides in the 58 L1 rebuilt sequence are indicated with dots.

FIGURE 2 shows the rebuilt synthetic HPV 58 L1 double-stranded nucleic acid and single-code amino acid sequence above. Nucleotide number is indicated to the left.

FIGURE 3 shows a Northern blot of HPV 58 L1 wt and 58 L1 R transcripts (see EXAMPLE 4). The blot was probed with a cocktail of equal amounts of DIG-labeled 58L1 wt and 58 L1 R DNA probes. The quantity of total RNA electrophoresed per lane is indicated. The arrow on the right indicates the predicted size of a full length 58 L1 transcript.

FIGURE 4 shows a Western Blot of HPV 58 L1 wt (58), and 58 L1 R (58R) proteins. HPV 16 L1 was included as a reference (16). Ten, five and two and one-half micrograms of total yeast protein extract were denatured and applied to a 10% SDS-PAGE gel. HPV 58 L1 protein was detected using a yeast-absorbed anti-trpE-HPV 31 L1 goat polyclonal antiserum which cross-reacts with 58 L1 and 16 L1. Molecular weight markers are indicted in kDa on the left

FIGURE 5 depicts the amount (ng) of intact HPV 58 L1 VLPs per microgram of total yeast protein captured and detected in an ELISA (see EXAMPLE 7). The results of two experiments conducted in duplicate are included. VLP expression of HPV 58 L1 wt (black and gray boxes) was 36 ng/ μ g total yeast protein. VLP expression of HPV 58

L1 R (white and hatched boxes), the rebuilt yeast-codon optimized 58 L1, was ~2-3 fold higher than HPV 58 L1 wt expression reaching 95 ng/μg total yeast protein in experiment #2.

FIGURE 6 shows a representative sample of HPV 58 VLPs composed of HPV 58 L1 R protein molecules, described herein, as visualized by transmission electron
5 microscopy (see EXAMPLE 8). The bar represents approximately 100 nm.

DETAILED DESCRIPTION OF THE INVENTION

The majority of cervical carcinomas are associated with infections of specific oncogenic types of human papillomavirus (HPV). The present invention relates to
10 compositions and methods to elicit or enhance immunity to the protein products expressed by genes of oncogenic HPV types. Specifically, the present invention provides polynucleotides encoding HPV58 L1, wherein the polynucleotides are codon-optimized for high-level expression in yeast. The present invention also provides HPV58 virus-like particles (VLPs), which are produced in yeast, and discloses use of said polynucleotides and VLPs in
15 pharmaceutical compositions and vaccines for the prevention and/or treatment of HPV-associated cancer.

A wild-type HPV58 L1 nucleotide sequence has been reported (Genbank Accession # NC_001443, *see* Kirii et al. *Virology* 185(1): 424-427 (1991)). The present invention provides synthetic DNA molecules encoding the HPV58 L1 protein. The
20 synthetic molecules of the present invention comprise a sequence of codons, wherein at least some of the codons have been altered to use the codons preferred by a yeast cell for high-level expression. The synthetic molecules may be used as a coding sequence for expression of HPV58 L1 protein, which may self-assemble into VLPs. Said VLPs may be used in a VLP-based vaccine to provide effective immunoprophylaxis against papillomavirus infection
25 through neutralizing antibody and cell-mediated immunity. Such VLP-based vaccines are also useful for treatment of already established HPV infections.

Expression of HPV VLPs in yeast cells offers the advantages of being cost-effective and easily adapted to large-scale growth in fermenters. In addition, the yeast genome can be readily altered to ensure selection of recombinant, transformed yeast with
30 increased growth and expression potential. However, many HPV L1 proteins, including HPV58 L1 are expressed at levels in yeast cells which are lower than what is desirable for commercial scale-up.

Accordingly, the present invention relates to HPV58 L1 gene sequences that are "optimized" for high-level expression in a yeast cellular environment.

A "triplet" codon of four possible nucleotide bases can exist in over 60 variant forms. Because these codons provide the message for only 20 different amino acids (as well as transcription initiation and termination), some amino acids can be coded for by more than one codon, a phenomenon known as codon redundancy. For reasons not
5 completely understood, alternative codons are not uniformly present in the endogenous DNA of differing types of cells. Indeed, there appears to exist a variable natural hierarchy or "preference" for certain codons in certain types of cells. As one example, the amino acid leucine is specified by any of six DNA codons including CTA, CTC, CTG, CTT, TTA, and TTG. Exhaustive analysis of genome codon use frequencies for microorganisms has
10 revealed endogenous DNA of *E. coli* most commonly contains the CTG leucine-specifying codon, while the DNA of yeasts and slime molds most commonly includes a TTA leucine-specifying codon. In view of this hierarchy, it is generally believed that the likelihood of obtaining high levels of expression of a leucine-rich polypeptide by an *E. coli* host will depend to some extent on the frequency of codon use. For example, it is likely that a gene
15 rich in TTA codons will be poorly expressed in *E. coli*, whereas a CTG rich gene will probably be highly expressed in this host. Similarly, a preferred codon for expression of a leucine-rich polypeptide in yeast host cells would be TTA.

The implications of codon preference phenomena on recombinant DNA techniques are manifest, and the phenomenon may serve to explain many prior failures to
20 achieve high expression levels of exogenous genes in successfully transformed host organisms—a less "preferred" codon may be repeatedly present in the inserted gene and the host cell machinery for expression may not operate as efficiently. This phenomenon suggests that synthetic genes which have been designed to include a projected host cell's preferred codons provide an optimal form of foreign genetic material for practice of recombinant
25 protein expression. Thus, one aspect of this invention is an HPV58 L1 gene that is codon-optimized for high-level expression in a yeast cell. In a preferred embodiment of this invention, it has been found that the use of alternative codons encoding the same protein sequence may remove the constraints on expression of HPV58 L1 proteins by yeast cells.

In accordance with this invention, HPV58 L1 gene segments were converted
30 to sequences having identical translated sequences but with alternative codon usage as described by Sharp and Cowe (Synonymous Codon Usage in *Saccharomyces cerevisiae*. *Yeast* 7: 657-678 (1991)), which is hereby incorporated by reference. The methodology generally consists of identifying codons in the wild-type sequence that are not commonly associated with highly expressed yeast genes and replacing them with optimal codons for
35 high expression in yeast cells. The new gene sequence is then inspected for undesired

sequences generated by these codon replacements (e.g., "ATTTA" sequences, inadvertent creation of intron splice recognition sites, unwanted restriction enzyme sites, high GC content, presence of transcription termination signals that are recognized by yeast, etc.). Undesirable sequences are eliminated by substitution of the existing codons with different
5 codons coding for the same amino acid. The synthetic gene segments are then tested for improved expression.

The methods described above were used to create synthetic gene segments for HPV58 L1, resulting in a gene comprising codons optimized for high-level expression. While the above procedure provides a summary of our methodology for designing codon-
10 optimized genes for use in HPV vaccines, it is understood by one skilled in the art that similar vaccine efficacy or increased expression of genes may be achieved by minor variations in the procedure or by minor variations in the sequence.

Accordingly, the present invention relates to a synthetic polynucleotide comprising a sequence of nucleotides encoding an HPV58 L1 protein, or a biologically active
15 fragment or mutant form of an HPV58 L1 protein, the polynucleotide sequence comprising codons optimized for expression in a yeast host cell. Said mutant forms of the HPV58 L1 protein include, but are not limited to: conservative amino acid substitutions, amino-terminal truncations, carboxy-terminal truncations, deletions, or additions. Any such biologically active fragment and/or mutant will encode either a protein or protein fragment which at least
20 substantially mimics the immunological properties of the HPV58 L1 protein as set forth in SEQ ID NO:2. The synthetic polynucleotides of the present invention encode mRNA molecules that express a functional HPV58 L1 protein so as to be useful in the development of a therapeutic or prophylactic HPV vaccine.

One aspect of this invention is a codon-optimized nucleic acid molecule
25 which encodes the HPV58 L1 protein as set forth in SEQ ID NO:2, said nucleic acid molecule comprising a sequence of nucleotides as set forth in SEQ ID NO:1.

The present invention also relates to recombinant vectors and recombinant host cells, both prokaryotic and eukaryotic, which contain the nucleic acid molecules disclosed throughout this specification. In a preferred embodiment of this invention, the host
30 cell is a yeast host cell.

The synthetic HPV58 DNA or fragments thereof constructed through the methods described herein may be recombinantly expressed by molecular cloning into an expression vector containing a suitable promoter and other appropriate transcription regulatory elements, and transferred into prokaryotic or eukaryotic host cells to produce
35 recombinant HPV58 L1. Techniques for such manipulations are fully described by

Sambrook et al. (*Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989); *Current Protocols in Molecular Biology*, Ausubel et al., Green Pub. Associates and Wiley-Interscience, New York (1988); *Yeast Genetics: A Laboratory Course Manual*, Rose et al., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1990)), which are hereby incorporated by reference in their entirety.

Thus, the present invention relates to a process for expressing an HPV58 L1 protein in a recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid encoding an HPV58 L1 protein into a yeast host cell; and (b) culturing the yeast host cell under conditions which allow expression of said HPV58 L1 protein.

The present invention further relates to a process for expressing an HPV58 L1 protein in a recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid encoding an HPV58 L1 protein into a yeast host cell; wherein the nucleic acid molecule is codon-optimized for optimal expression in the yeast host cell and; (b) culturing the yeast host cell under conditions which allow expression of said HPV58 L1 protein.

This invention further relates to a process for expressing an HPV58 L1 protein in a recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid as set forth in SEQ ID NO:1 into a yeast host cell; and, (b) culturing the yeast host cell under conditions which allow expression of said HPV58 L1 protein.

The synthetic genes of the present invention can be assembled into an expression cassette that comprises sequences designed to provide efficient expression of the HPV58 L1 protein in the host cell. The cassette preferably contains the synthetic gene, with related transcriptional and translations control sequences operatively linked to it, such as a promoter, and termination sequences. In a preferred embodiment, the promoter is the *S. cerevisiae GAL1* promoter, although those skilled in the art will recognize that any of a number of other known yeast promoters such as the *GAL10*, *GAL7*, *ADH1*, *TDH3* or *PGK* promoters, or other eukaryotic gene promoters may be used. A preferred transcriptional terminator is the *S. cerevisiae ADH1* terminator, although other known transcriptional terminators may also be used. The combination of *GAL1* promoter – *ADH1* terminator is particularly preferred.

This invention further provides an isolated and purified HPV 58 L1 polypeptide comprising a sequence of amino acids as set forth in SEQ ID NO:2.

Another aspect of this invention is an HPV58 virus-like particle (VLP) produced by recombinantly expressing the HPV58 L1 or L1 + L2 genes in a yeast cell, methods of producing HPV58 VLPs, and methods of using HPV58 VLPs. VLPs can self-assemble when L1, the major capsid protein of human and animal papillomaviruses, is

expressed in yeast, insect cells, mammalian cells or bacteria (for review, *see* Schiller and Roden, in *Papillomavirus Reviews: Current Research on Papillomaviruses*; Lacey, ed. Leeds, UK: Leeds Medical Information, pp 101-12 (1996)). Morphologically indistinct HPV VLPs can also be produced by expressing a combination of the L1 and L2 capsid proteins.

5 VLPs are composed of 72 pentamers of L1 in a T=7 icosahedral structure (Baker et al., *Biophys. J.* 60(6): 1445-56 (1991)).

VLPs are morphologically similar to authentic virions and are capable of inducing high titres of neutralizing antibodies upon administration into an animal. Immunization of rabbits (Breitburd et al., *J. Virol.* 69(6): 3959-63 (1995)) and dogs (Suzich et al., *Proc. Natl. Acad. Sci. USA* 92(25): 11553-57 (1995)) with VLPs was shown to both

10 induce neutralizing antibodies and protect against experimental papillomavirus infection. However, because the VLPs do not contain the potentially oncogenic viral genome and can self-assemble when expressed from a single gene, they present a safe alternative to the use of live virus in HPV vaccine development (for review, *see* Schiller and Hidesheim, *J. Clin.*

15 *Virol.* 19: 67-74 (2000)).

Thus, the present invention relates to virus-like particles comprised of recombinant L1 protein or recombinant L1 + L2 proteins of HPV58, wherein the recombinant protein is expressed in a yeast cell.

As stated above, in a preferred embodiment of the invention, the HPV58

20 VLPs are produced in yeast. In a further preferred embodiment, the yeast is selected from the group consisting of: *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Pichia pastoris*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, and *Schizosaccharomyces pombe*.

Another aspect of this invention is an HPV58 VLP which comprises an HPV58 L1 protein produced by a codon-optimized HPV58 L1 gene. In a preferred

25 embodiment of this aspect of the invention, the codon-optimized HPV58 L1 gene comprises a sequence of nucleotides as set forth in SEQ ID NO:1.

Yet another aspect of this invention is a method of producing HPV58 VLPs, comprising: (a) transforming yeast with a recombinant DNA molecule encoding HPV58 L1 protein or HPV58 L1 + L2 proteins; (b) cultivating the transformed yeast under conditions

30 that permit expression of the recombinant DNA molecule to produce the recombinant HPV58 protein; and (c) isolating the recombinant HPV58 protein to produce HPV58 VLPs.

In a preferred embodiment of this aspect of the invention, the yeast is transformed with a codon-optimized HPV58 L1 gene to produce HPV58 VLPs. In a particularly preferred embodiment, the codon-optimized HPV58 L1 gene comprises a

35 sequence of nucleotides as set forth in SEQ ID NO:1.

This invention also provides a method for inducing an immune response in an animal comprising administering HPV58 virus-like particles to the animal. In a preferred embodiment, the HPV58 VLPs are produced by a codon-optimized gene.

Yet another aspect of this invention is a method of preventing and/or treating HPV-associated cervical cancer comprising administering to a mammal a vaccine comprising HPV58 VLPs. In a preferred embodiment of this aspect of the invention, the HPV58 VLPs are produced in yeast.

This invention also relates to a vaccine comprising HPV58 virus-like particles (VLPs).

In an alternative embodiment of this aspect of the invention, the vaccine further comprises VLPs of at least one additional HPV type. In a preferred embodiment, the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV45, HPV51, HPV52, HPV55, HPV56, HPV59, and HPV68.

In a preferred embodiment of this aspect of the invention, the vaccine further comprises HPV16 VLPs.

In another preferred embodiment of the invention, the vaccine further comprises HPV16 VLPs and HPV18 VLPs.

In yet another preferred embodiment of the invention, the vaccine further comprises HPV6 VLPs, HPV11 VLPs, HPV16 VLPs and HPV18 VLPs.

This invention also relates to pharmaceutical compositions comprising HPV 58 virus-like particles. Further, this invention relates to pharmaceutical compositions comprising HPV58 VLPs and VLPs of at least one additional HPV type. In a preferred embodiment, the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV45, HPV51, HPV52, HPV55, HPV56, HPV59, and HPV68.

Vaccine compositions of the present invention may be used alone at appropriate dosages which allow for optimal inhibition of HPV58 infection with minimal potential toxicity. In addition, co-administration or sequential administration of other agents may be desirable.

The amount of virus-like particles to be introduced into a vaccine recipient will depend on the immunogenicity of the expressed gene product. In general, an immunologically or prophylactically effective dose of about 10 µg to 100 µg, and preferably about 20 µg to 60 µg of VLPs is administered directly into muscle tissue. Subcutaneous injection, intradermal introduction, impression through the skin, and other modes of

administration such as intraperitoneal, intravenous, or inhalation delivery are also contemplated. It is also contemplated that booster vaccinations may be provided. Parenteral administration, such as intravenous, intramuscular, subcutaneous or other means of administration with adjuvants such as alum or Merck aluminum adjuvant, concurrently with
5 or subsequent to parenteral introduction of the vaccine of this invention is also advantageous.

All publications mentioned herein are incorporated by reference for the purpose of describing and disclosing methodologies and materials that might be used in connection with the present invention. Nothing herein is to be construed as an admission that
10 the invention is not entitled to antedate such disclosure by virtue of prior invention.

Having described preferred embodiments of the invention with reference to the accompanying drawings, it is to be understood that the invention is not limited to those precise embodiments, and that various changes and modifications may be effected therein by one skilled in the art without departing from the scope or spirit of the invention as defined in
15 the appended claims.

The following examples illustrate, but do not limit the invention.

EXAMPLE 1

Determination of a Representative HPV 58 L1 Sequence

20 The HPV 58 L1 sequence has been described previously (Genbank Accession # NC_001443). It is not uncommon, however, to find minor sequence variations between DNAs obtained from clinical isolates. To determine a representative HPV58 L1 wild-type sequence, DNA was isolated from three clinical samples previously shown to contain HPV 58 DNA. HPV 58 L1 sequences were amplified in a polymerase chain reaction
25 (PCR) using *Taq* DNA polymerase and the following primers: HPV 58 L1 F 5' – A T G T C C G T G T G G C G G C C T A G T – 3' (SEQ ID NO:4) and 58 3' I1 BglII 5' – G A G A T C T G T G T A A G T A C C A C A A C A A T T A – 3' (SEQ ID NO:5). The amplified products were electrophoresed on agarose gels and visualized by ethidium bromide staining. The ~ 1500 bp L1 bands were excised and DNA purified using GeneClean Spin Kit (Q-Bio
30 Gene, Carlsbad, CA). The DNA was then ligated to the TA cloning vector, pCR2.1 (Invitrogen). TOP10F' *E. Coli* were transformed with the ligation mixture and plated on LB agar with ampicillin plus IPTG and X-gal for blue/white colony selection. The plates were inverted and incubated for 16 hours at 37°C. White colonies were cultured in LB medium with ampicillin by shaking at 37°C for 16 hours. Minipreps were performed to extract the
35 plasmid DNA.

To demonstrate the presence of the L1 gene in the plasmids, restriction endonuclease digestions were conducted. Restriction fragments were viewed by agarose gel electrophoresis and ethidium bromide staining. DNA sequencing was performed on plasmids containing cloned L1 inserts from each of the three clinical isolates. In order to generate a reference sequence for later optimization, the nucleotide and translated amino acid sequences from each of the clones were compared to the published HPV 58 L1 sequences. Sequence analysis of the three clinical isolates revealed that no sequence was identical to the Genbank sequence. The pCR2.1 HPV 58L1 clone #4 was chosen to be the representative 58 L1 sequence and is referred to interchangeably herein as the "58 L1 wild-type sequence" or "58 wt sequence" (SEQ ID NO:3, see FIGURE 1). The 58 L1 wt sequence contained five point mutations: two resulting in amino acids changes and three silent point mutations. The point mutations resulting in amino acid changes with respect to the Genbank HPV 58 L1 sequence were located at nucleotide 372 (A->T), altering amino acid 124 from leucine to phenylalanine, and nucleotide 897 (A->G), altering amino acid 299 from isoleucine to methionine. The three silent point mutations were located at nucleotides 774 (A->G), 792 (T->C), and 999 (G->A).

The 58 L1 wild-type sequence was PCR-amplified using *Taq* polymerase and the following primers, which add *Bam*HI extensions: 5'58 BamHI 5' – G G G A T C C C A C A A A A C A A A A T G T C C G T G T G G C – 3' (SEQ ID NO:6) and 3' Bam 58 5' – G G G A T C C G T G T A A G T A C C A C A A C A A T T A – 3' (SEQ ID NO: 7). The resulting PCR products were visualized by agarose gel electrophoresis, followed by ethidium bromide staining. The ~ 1500 bp band was excised and DNA-purified using the GeneClean kit. The PCR product was then ligated to pCR2.1 and TOP10F' cells were transformed with the ligation mixture. White colonies were selected and cultured in LB medium with ampicillin by shaking at 37°C for 16 hours. Minipreps were performed to extract the plasmid DNA. To release the HPV 58 L1 gene from the vector sequences, *Bam*HI restriction endonuclease digestions were performed. The digested DNA was subjected to agarose gel electrophoresis and viewed by ethidium bromide staining. The L1 band was purified using the GeneClean kit and ligated to dephosphorylated, *Bam*HI digested pGAL110. DH5α *E.coli* cells were transformed with the ligation mixture. To screen for the HPV 58 L1 insert in the correct orientation, plasmid DNA from colonies was PCR-amplified. DNA sequencing was conducted to confirm sequence and orientation of the inserts. A single clone was selected and named pGAL110-HPV 58L1 #10. Maxiprep DNA from the selected clone was prepared. *Saccharomyces cerevisiae* cells were made competent by spheroplasting with glucylase and transformed with pGAL110-HPV 58L1 #1. The yeast transformation mixture was plated in

Leu(-) sorbitol top-agar onto Leu(-) sorbitol agar plates and incubated inverted for 3-5 days at 30°C. Colonies were picked and streaked for isolation on Leu(-) sorbitol agar plates. Isolated colonies were subsequently grown in 5 ml of 5 X Leu(-) Ade(-) sorbitol with 1.6% glucose and 4% galactose in rotating tube cultures at 30°C to induce 58 L1 transcription and protein expression.

EXAMPLE 2

Yeast codon optimization

Yeast-preferred codons have been described (Sharp, Paul M and Cowe, Elizabeth. Synonymous Codon Usage in *Saccharomyces cerevisiae* YEAST 7: 657-678 (1991)). Expression of the HPV 58 L1 wt protein was detectable, however to obtain increased expression, the HPV 58 L1 gene was rebuilt utilizing the preferred yeast codons. The rebuilt 58 L1 sequence, which comprises yeast-optimized codon sequences, contained 404 nucleotide alterations compared to the 58 L1 wt sequence. The resulting sequence is referred to herein as "58 L1 R" (R = rebuild, see FIGURE 1). The translated amino acid sequence of 58 L1 R was not altered. The nucleotide (SEQ ID NO:1) and amino acid (SEQ ID NO:2) sequences of HPV 58 L1 R are shown in FIGURE 2. Said rebuilt sequence provides increased HPV 58 L1 expression, which is a significant advance over the wild-type for use in vaccine development (see EXAMPLE 4).

The strategy employed to produce the optimized gene was to design long overlapping sense and antisense oligomers that span the gene, substituting nucleotides with yeast-preferred codon sequences, while maintaining the amino acid sequence. These oligomers were used in place of template DNA in a PCR reaction with *Pfu* polymerase. Additional amplification primers were designed and used to amplify the rebuilt sequences from template oligomers.

The optimal conditions for amplification were section-specific, however, most employed a program resembling 95°C for 2 minutes (denaturing) followed by 35 cycles of 95°C for 1 minute (denaturing), 55°C for 1 minute (annealing), 72°C for 3.5 minute (extension), followed by a 72°C for 10 minute final extension and 4°C hold. PCR products were examined by agarose gel electrophoresis. Bands of the appropriate size were excised and DNA gel purified. The amplified fragments were then used as templates to assemble the 1497 nt rebuilt HPV 58 L1 gene.

Following rebuild, the 1497 nt band was gel purified and ligated to pCR-Blunt vector (Invitrogen, Carlsbad, CA). Following ligation, TOP10 cells were transformed with the ligation mixture. Colonies were grown in LB with kanamycin and plasmid DNA

was extracted from the colonies by miniprep techniques. The plasmid DNA was sequenced to confirm the desired 58 L1 rebuild changes. To add *Bam*HI extensions to both ends, the 58 L1 R (rebuild) was re-amplified from pCR-Blunt-58 L1 R with the following primers: 5'Bam 58 Rebuild 5' – G G A T C C C A C A A A C A A A A T G T C T G T C T G G A G A C C – 3' (SEQ ID NO:8) and 3'Bam 58 Rebuild 5' – G G A T C C T T A C T T C T T G A C C T T C – 3' (SEQ ID NO:9).

The amplified L1 product was gel-purified using the GeneClean kit and cloned into pCR2.1 (Invitrogen). Top10F' cells were transformed with the pCR2.1 plasmid. White colonies were cultured in LB medium with ampicillin, shaking at 37°C for 16 hours. Minipreps were performed to extract the plasmid DNA. To release the HPV 58 L1 gene from the vector sequences, *Bam*HI restriction endonuclease digestions were performed. The digested DNA was subjected to agarose gel electrophoresis and viewed by ethidium bromide staining. The L1 band was purified using the GeneClean kit and ligated to dephosphorylated, *Bam*HI-digested pGAL110. DH5α *E.coli* cells were transformed with the ligation mixture. The resulting colonies were screened by PCR for the HPV 58 L1 insert in the correct orientation. Maxiprep DNA was prepared. Sequence and orientation were confirmed by restriction digest profiles and DNA sequencing. The selected clone was named pGAL110-HPV 58L1R #17. *Saccharomyces cerevisiae* cells were made competent by spheroplasting and transformed with pGAL110-HPV 58L1R #17. The yeast transformation was plated in Leu(-) sorbitol top-agar on Leu(-) sorbitol agar plates and incubated inverted for 3-5 days at 30°C. Colonies were picked and streaked for clonal isolation on Leu(-) sorbitol agar plates. Isolated colonies were subsequently grown in 5 ml of 5 X Leu(-) Ade(-) sorbitol with 1.6% glucose and 4% galactose in rotating tube cultures at 30°C to induce L1 transcription and protein expression. After 48 hours, a culture volume equivalent to an OD₆₀₀ = 10 amount of cells was pelleted, the supernatant was removed and the pellets were frozen and stored at -70°C.

EXAMPLE 3

RNA preparation

Cell pellets of transformed yeast cells induced to express HPV 58 L1 or HPV 58 L1 R by galactose induction were thawed on ice, suspended in 0.8 ml of Trizol reagent (Life Technologies, Gibco BRL) and incubated at room temperature for 5 minutes. One fifth volume of chloroform was added to the vial. It was then shaken vigorously for 15 seconds to mix and incubated at room temperature for 3 minutes. After a 5 minute centrifugation at 13 k rpms, the upper phase was collected and transferred to a new vial. 0.4 ml isopropanol was

added and incubated at room temperature for 10 minutes. To pellet the RNA, centrifugation was performed at 13 k rpms for 10 minutes. The supernatant was decanted, the RNA pellet washed with 75% EtOH and centrifugation repeated. The supernatant was decanted and the RNA pellet allowed to air dry for 15 minutes followed by suspension in RNase-free water.

- 5 Spectrophotometry was performed to determine the concentration of RNA in the sample using the assumption that an A₂₆₀ reading of 1 = 40 µg/ml RNA when the A₂₆₀/A₂₈₀ is 1.7-2.0.

EXAMPLE 4

10 Northern Blot Analysis

- A 1.1% agarose formaldehyde gel was cast. Five and ten micrograms of RNA were combined with denaturing buffer (final concentrations: 6% formaldehyde, 50% formamide and 0.1 x MOPS) and heated to 65°C for 10 minutes. A one-tenth volume of gel loading buffer was added and the sample loaded onto the gel. Electrophoresis was performed
15 at 75 volts in 1 x MOPS buffer for ~ 3 hours. The gel was washed for 60 minutes in 10 x SSC.

- The RNA was transferred to a Hybond-N+ nylon membrane (Amersham Biosciences, Piscataway, NJ) by capillary action over 16 hours in 10 x SSC. The RNA was then fixed to the nylon membrane by cross-linking using the Stratagene UV Stratalinker auto-crosslink function (Stratagene, La Jolla, CA). After fixing, the nylon membrane was allowed
20 to air dry.

- The Roche DIG High Prime DNA Labeling and Detection Kit I (Hoffmann-La Roche Ltd., Basel, Switzerland) was used to label 58 L1 wt and 58 L1 R DNA sequences with DIG to be used as a probe cocktail to detect 58 L1 wt and 58 L1 R transcripts on the
25 Northern blot. The pre-hybridization, hybridization and immunological development using an anti-DIG alkaline phosphatase conjugated antibody were performed per the manufacturer's recommendations. Briefly, the blot was pre-hybridized at 37°C for 30 minutes with gentle shaking. The probe cocktail was denatured by heating to 95°C for 5 minutes and quenching on ice. The probe cocktail was added to the hybridization solution
30 and applied to the membrane for 4 hours at 44.6°C with gentle shaking. The hybridization solution was then removed and the blot washed 2 x for 5 minutes in 2 x SSC with 0.1% SDS at room temperature, followed by an additional wash at 65°C with 0.5 x SSC and 0.1% SDS. The blot was then blocked for 30 minutes and anti-DIG alkaline phosphatase conjugated antibody was applied at a 1:5000 dilution for 30 minutes. The blot was washed and the

presence of probe-bound RNA was determined by NBT/BCIP substrate detection of the alkaline phosphatase conjugated anti-DIG bound antibody.

Initial analysis of yeast expressing 58 L1 wt suggested that there was functional HPV 58 L1 full-length transcription and translation; however, the level of expression might be increased if the sequence was rebuilt with yeast-preferred codon sequences. The rebuilt 58 L1 sequence was engineered to omit any possible premature transcription termination sites to ensure robust transcription. Northern blot analysis of the 58 L1 R transcript revealed that increased amounts of full-length transcripts were generated compared to the results seen for 58 L1 wt (FIGURE 3).

10

EXAMPLE 5

HPV 58 L1 Protein Expression

Frozen yeast cell pellets of galactose-induced cultures equivalent to an OD₆₀₀= 10 quantity of cells, were thawed on ice and suspended in 300 µl of PC buffer (100 mM Na₂HPO₄ and 0.5 M NaCl, pH 7.0) with 2mM PMSF. Acid-washed 0.5mm glass beads were added at a concentration of ~ 0.5g/tube. The tubes were vortexed for 3 cycles of 5 minutes at 4°C with a 1 minute break. 7.5 µl of 20% TritonX100 was added and the vortex step was repeated for 5 minutes at 4°C. The tubes were placed on ice for 15 minutes, followed by centrifugation for 10 minutes at 4°C. The supernate was transferred to a sterile microfuge tube, labeled as total yeast protein extract, dated and stored at -70°C.

20

EXAMPLE 6

Western Blot Analysis

Total yeast protein extract from twenty isolated yeast colonies for each 58 L1 construct were analyzed by Western blot to confirm expression of 58 L1 protein after galactose induction.

25

Ten, five and two and one-half micrograms of total yeast protein extract of representative 58 L1 wt and 58 L1 R isolates were combined with SDS-PAGE loading buffer and heated to 95°C for 10 minutes. The 16 L1 protein, which is approximately 55 kD, was included as a positive control, along with HPV L1-free total yeast protein extract as a negative control (data not shown).

30

The proteins were loaded onto a 10% SDS-PAGE gel and electrophoresed in Tris-Glycine buffer. After protein separation, the proteins were Western transferred from the gel to nitrocellulose and the blot was blocked in 1 x diluent buffer (Kirkegaard and Perry Laboratories, Gaithersburg, MD) for 1 hour at room temperature with rocking. The blot was

35

washed three times and incubated at room temperature for 16 hours with yeast absorbed goat anti-trpE-HPV 31 L1 serum, which cross-reacts with HPV 16 and HPV 58 L1 proteins. The blot was then washed three times and incubated with a 1:2500 dilution of anti-goat-HRP conjugated antibody for 1 hr. The blot was again washed three times and NBT/BCIP
5 detection substrate applied (Kirkegaard and Perry Laboratories). Immunoreactive proteins were detected as purple bands on the blot.

In all cases, the 58 L1 protein was detected as a distinct immunoreactive band on the nitrocellulose corresponding to approximately 55 kD (FIGURE 4). The intensity of the 58 L1 R bands appeared to be greater than that seen for 58 L1 wt, suggesting
10 improved 58 L1 expression was achieved by yeast-codon optimization.

EXAMPLE 7

ELISA Assay

To demonstrate 58 L1 VLP expression, a portion of 58 L1 wt and 58 L1 R
15 total yeast protein extract was analyzed by ELISA. The yeast cells expressing HPV 58 L1 and HPV 58 L1 R were grown by a variety of methods, including rotating tube cultures, shake flasks, and fermenters. The yeast cells were lysed and protein extracts were made to determine the amount of HPV 58 L1 virus-like particles (VLPs) produced per microgram of total protein. A sandwich ELISA was designed to demonstrate HPV 58 L1 VLP expression.

20 Protein G purified H582C3.F7 (F7) monoclonal antibody (mAb) was used to bind intact 58 L1 VLPs found in the yeast protein extracts. F7 specifically recognizes an HPV 58 L1 VLP conformational epitope. The unbound proteins were washed away and H586E11.F4 (F4), another HPV 58 L1 VLP conformational specific mAb, was applied as a detection antibody. True, conformationally correct, 58 L1 VLPs were bound and detection
25 was facilitated by the use of an anti-mouse IgG2b HRP-conjugated antibody and TMB substrate.

Specifically, F7 was used to coat the bottom of Immulon 4 HBX 96 well plates overnight at 4°C. The plates were washed three times with PBS and 0.05% Tween 20, followed by blocking with blocking solution (PBS + 0.05% Tween 20 + 1% BSA). The
30 plates were washed three times and antigens (total yeast cell lysates diluted in blocking solution to 12.5 µg/ml) were applied to row A in duplicate. Reference standards of purified HPV 58 L1 VLPs were applied to row A columns 3 and 4 at 206ng/ml in 12.5 µg/ml total yeast protein. The reference and test samples were then serially diluted two-fold down each column. After three hours at room temperature, the excess antigen was removed by
35 aspiration and the plates were washed 3 times. F4 conformational specific mAb was diluted

in blocking solution and applied to each well for one hour at room temperature. The plates were washed three times and an anti-mouse IgG2b HRP-conjugated antibody was diluted in blocking solution and applied for 1 hour at room temperature. The plates were washed and TMB (Pierce Biotechnology, Inc., Rockford, IL) was applied for 5 minutes to detect HRP-conjugated antibody complexes. The detection reaction was stopped with the addition of 2M H₂SO₄. Plates were read at 450 nm wavelength and the concentration of HPV 58 L1 VLP was determined by comparison to the reference standards in ng VLP / μ g total protein.

FIGURE 5 shows a comparison of the amount of VLPs detected/ μ g of total protein from yeast expressing HPV 58 L1 wt and HPV 58 L1 R from two separate experiments. HPV 58 L1 VLP expression levels increased ~2-3 fold with yeast-codon optimization

EXAMPLE 8

Transmission Electron Microscopy

To demonstrate that the 58 L1 protein was in fact self-assembling to form pentameric-L1 capsomers, which in turn self-assemble into virus-like particles, a partially purified 58 L1 R protein extract was subjected to transmission electron microscopy (EM).

Yeast were grown under small scale fermentation and pelleted. The resulting pellets were subjected to purification treatments. Pellet and clarified yeast extracts were analyzed by immunoblot to demonstrate L1 protein expression and retention through the purification procedure. Clarified yeast extracts were then subjected to centrifugation over a 45%-sucrose cushion and the resulting pellet suspended in buffer for analysis by transmission EM.

A representative image of the 58 L1 R VLPs produced is shown in FIGURE 6. The diameter of the spherical particles in this crude sample ranged from 30 to 60 nm, with some particles displaying a regular array of capsomers.

WHAT IS CLAIMED IS:

1. A nucleic acid molecule comprising a sequence of nucleotides that encodes an HPV58 L1 protein as set forth in SEQ ID NO:2, the nucleic acid sequence being
5 codon-optimized for high-level expression in a yeast cell.
2. A vector comprising the nucleic acid molecule of claim 1.
3. A host cell comprising the vector of claim 2.
10
4. The host cell of claim 3, wherein the host cell is a yeast cell.
5. The host cell of claim 4, wherein the yeast cell is selected from the group consisting of: *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Pichia pastoris*,
15 *Kluyveromyces fragilis*, *Kluyveromyces lactis*, and *Schizosaccharomyces pombe*.
6. The host cell of claim 5, wherein the host cell is *Saccharomyces cerevisiae*.
7. The nucleic acid molecule of claim 1, wherein the sequence of
20 nucleotides comprises a sequence of nucleotides as set forth in SEQ ID NO:1.
8. A vector comprising the nucleic acid molecule of claim 7.
9. A host cell comprising the vector of claim 8.
25
10. Virus-like particles (VLPs) comprised of recombinant L1 protein or recombinant L1 + L2 proteins of HPV58, wherein the recombinant L1 protein or the recombinant L1 + L2 proteins are produced in yeast.
30
11. The VLPs of claim 10, wherein the recombinant L1 protein or recombinant L1 + L2 proteins are encoded by a codon-optimized HPV58 L1 nucleic acid molecule.

12. The VLPs of claim 11, wherein the codon-optimized nucleic acid molecule comprises a sequence of nucleotides as set forth in SEQ ID NO:1.

5 13. A method of producing the VLPs of Claim 11, comprising:
(a) transforming yeast with a codon-optimized DNA molecule encoding HPV58 L1 protein or HPV58 L1 + L2 proteins;
(b) cultivating the transformed yeast under conditions that permit expression of the codon-optimized DNA molecule to produce a recombinant papillomavirus protein; and
10 (c) isolating the recombinant papillomavirus protein to produce the VLPs of Claim 11.

14. A vaccine comprising the VLPs of Claim 11.

15 15. Pharmaceutical compositions comprising the VLPs of claim 11.

16. A method of preventing HPV infection comprising administering the vaccine of Claim 14 to a mammal.

20 17. A method for inducing an immune response in an animal comprising administering the VLPs of Claim 11 to an animal.

25 18. The virus-like particles of Claim 11 wherein the yeast is selected from the group consisting of *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Pichia pastoris*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, and *Schizosaccharomyces pombe*.

19. The virus-like particles of claim 18, wherein the yeast is *Saccharomyces cerevisiae*.

30 20. The vaccine of claim 14, further comprising VLPs of at least one additional HPV type.

21. The vaccine of claim 20 wherein the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV45, HPV51, HPV52, HPV55, HPV56, HPV59, and HPV68.
35

22. The vaccine of claim 21, wherein the at least one HPV type comprises HPV16.
- 5 23. The vaccine of claim 22, further comprising HPV18 VLPs.
24. The vaccine of claim 23, further comprising HPV6 VLPs and HPV11 VLPs.
- 10 25. The vaccine of claim 24, further comprising HPV31 VLPs.
26. The vaccine of claim 23, further comprising HPV31 VLPs.
27. The vaccine of claim 25, further comprising HPV45 VLPs.
- 15 28. The vaccine of claim 26, further comprising HPV45 VLPs.
29. An isolated and purified HPV 58 L1 polypeptide comprising a sequence of amino acids as set forth in SEQ ID NO:2.
- 20

1/10

HPV 58 L1 Nucleotide Sequence Alignment

58 L1 wt	(1)	ATGTCCGTGTGGCGGCCTAGTGAGGCCACTGTGTACCTGCCTCCTGTGCC
58 L1 R	(1)C...A.A..ATCC..A..T..C..C...T....A..A..T..
58 L1 wt	(51)	TGTGTCTAAGGTTGTAAGCACTGATGAATATGTGTCACGCACAAGCATT
58 L1 R	(51)	A..C..C.....C..CTC.....C.....C..C..TA.A..CTCT..C.
58 L1 wt	(101)	ATTATTATGCTGGCAGTTCAGACTTTTGGCTGTTGGCAATCCATATTTT
58 L1 R	(101)	.C..C..C.....TTCC..T...T.G.....T..C.....C..C
58 L1 wt	(151)	TCCATCAAAAGTCCCAATAACAATAAAAAAGTATTAGTTCCCAAGGTATC
58 L1 R	(151)GTC...A..C.....C..G..G..C..G.....A.....C..
58 L1 wt	(201)	AGGCTTACAGTATAGGGTCTTTAGGGTGCGTTTACCTGATCCCAATAAAT
58 L1 R	(201)	T..T..G..A..C..A.....C..A..CA.A..G..A..C..A..C..G.
58 L1 wt	(251)	TTGGTTTTCTGATACATCTTTTATAACCCTGATACACAACGTTTGGTC
58 L1 R	(251)	.C.....C..A..C..T..C..C..C.....A..C..T...A.A.....
58 L1 wt	(301)	TGGGCATGTGTAGGCCTTGAAATAGGTAGGGGACAGCCATTGGGTGTTGG
58 L1 R	(301)T.....C..TT.G.....C.....A..T..A.....
58 L1 wt	(351)	CGTAAGTGGTCATCCTTATTTCAATAAATTTGATGACACTGAAACCAGTA
58 L1 R	(351)	T..CTC.....C..A..C.....C..G..C..C.....C.....TCC.
58 L1 wt	(401)	ACAGATATCCCGCACAGCCAGGGTCTGATAACAGGGAATGCTTATCTATG
58 L1 R	(401)C..A..T..A.....T.....C.....A.....T..G..C...
58 L1 wt	(451)	GATTATAAACAAACACAATTATGTTTAATTGGCTGTAAACCTCCCACTGG
58 L1 R	(451)	..C..C..G.....C.....G.....G..C..T.....G..A..A.....
58 L1 wt	(501)	TGAGCATTGGGGTAAAGGTGTTGCCTGTAACAATAATGCAGCTGCTACTG
58 L1 R	(501)	...A..C.....G.....T.....C..C..T.....C.
58 L1 wt	(551)	ATTGTCCTCCATTGGAACCTTTTAATTCTATTATTGAGGATGGTGACATG
58 L1 R	(551)	.C.....A.....T.G..C..C..C..C..C..A..C.....
58 L1 wt	(601)	GTAGATACAGGGTTTGGATGCATGGACTTTGGTACATTGCAGGCTAATAA
58 L1 R	(601)	..C..C..T..T..C..T..T.....C.....C.....A.....C..

FIG.1A

2/10

58 L1 wt	(651)	AAGTGATGTGCCTATTGATATTTGTAACAGTACATGCAAATATCCAGATT
58 L1 R	(651)	GTCC..C..T..A..C..C..C.....TCC..C..T..G..C.....C.
58 L1 wt	(701)	ATTTAAAAATGGCCAGTGAACCTTATGGGGATAGTTTGTTCCTTTTTTCTT
58 L1 R	(701)	.C..G..G.....TTC.....A..C..T..CTCC.....C..CT.G
58 L1 wt	(751)	AGACGTGAGCAGATGTTTGTTAGGCACTTTTTTAATAGGGCCGGAAACT
58 L1 R	(751)	...A.A..A..A.....C..C..A.....C..C..C..A..T..T..GT.
58 L1 wt	(801)	TGGCGAGGCTGTCCCGGATGACCTTTATATTAAAGGGTCCGGTAATACTG
58 L1 R	(801)	G..T..A.....T..A..C...T.G..C..C..G..T..T.....C..C.
58 L1 wt	(851)	CAGTTATCCAAAGTAGTGCATTTTTTCCAACCTCCTAGTGGCTCTATGGTT
58 L1 R	(851)	.T..C.....TCCTC...T..C..C.....ATC...T..C.....C
58 L1 wt	(901)	ACCTCAGAATCACAATTATTTAATAAGCCTTATTGGCTACAGCGTGCACA
58 L1 R	(901)T.....T.....G..C..C.....A..C...T.G..AA.A..T..
58 L1 wt	(951)	AGGTCATAACAATGGCATTTGCTGGGGCAATCAGTTATTTGTTACCGTAG
58 L1 R	(951)C.....C..T..C..T.....T..C..A..G..C..C..T..C.
58 L1 wt	(1001)	TTGATACCACTCGTAGCACTAATATGACATTATGCACTGAAGTAACTAAG
58 L1 R	(1001)	.C..C.....A.ATC.....C.....C..G..T..C.....C..C...
58 L1 wt	(1051)	GAAGGTACATATAAAAATGATAATTTTAAGGAATATGTACGTCATGTTGA
58 L1 R	(1051)C..C..G..C..C..C..C.....C..CA.A..C..C..
58 L1 wt	(1101)	AGAATATGACTTACAGTTTGTTTTTCAGCTTTGCAAAATTACACTAACTG
58 L1 R	(1101)	G.....C.....G..A..C..C..C..AT.G..T..G..C..CT.G....
58 L1 wt	(1151)	CAGAGATAATGACATATATACATACTATGGATTCCAATATTTTGGAGGAC
58 L1 R	(1151)	.T..A..C.....C..C..C..C..C.....C..T..C..C.....A...
58 L1 wt	(1201)	TGGCAATTTGGTTTAACACCTCCTCCGTCTGCCAGTTTACAGGACACATA
58 L1 R	(1201)C.....G..T..A..A..A.....TTCC..G..A.....C..
58 L1 wt	(1251)	TAGATTTGTTACCTCCCAGGCTATTACTTGCCAAAAAACAGCACCCCTA
58 L1 R	(1251)	C.....C..C.....T..A.....C..C..T.....G..T..T..A..A.

FIG.1B

3/10

58 L1 wt	(1301)	AAGAAAAGGAAGATCCATTAAATAAATATACTTTTTGGGAGGTAACTTA
58 L1 R	(1301)	.G.....C.....G..C..G..C..C..C.....A..C.....G
58 L1 wt	(1351)	AAGGAAAAGTTTTCTGCAGATCTAGATCAGTTTCCTTTGGGACGAAAGTT
58 L1 R	(1351)C.....T..CT.G..C..A..C..A.....TA.....
58 L1 wt	(1401)	TTTATTACAATCAGGCCTTAAAGCAAAGCCCAGACTAAAACGTTTCGGCCC
58 L1 R	(1401)	C..G..G.....T..TT.G..G..T.....A...T.G..GA.A..T..T.
58 L1 wt	(1451)	CTACTACCCGTGCACCATCCACCAAACGCAAAAAGGTAAAAAATAA (SEQ ID NO:3)
58 L1 R	(1451)	.A..C..TA.A..T.....GA.A..G.....C..G..G (SEQ ID NO:1)

FIG.1C

4/10

Synthetic HPV 58 L1 Nucleotide and Amino Acid Sequences.

	M	S	V	W	R	P	S	E	A	T	V	Y	L	P	P	V	P
1	ATGTCCGTCT	GGAGACCATC	CGAAGCTACC	GTCTACTTGC	CACCAGTTCC												
	TACAGGCAGA	CCTCTGGTAG	GCTTCGATGG	CAGATGAACG	GTGGTCAAGG												
	V	S	K	V	V	S	T	D	E	Y	V	S	R	T	S	I	Y
51	AGTCTCCAAG	GTCGTCTCCA	CTGACGAATA	CGTCTCTAGA	ACCTCTATCT												
	TCAGAGGTTC	CAGCAGAGGT	GACTGCTTAT	GCAGAGATCT	TGGAGATAGA												
	Y	Y	A	G	S	S	R	L	L	A	V	G	N	P	Y	F	
101	ACTACTACGC	TGGTTCCTCT	AGATTGTTGG	CTGTTGGTAA	CCCATACTTC												
	TGATGATGCG	ACCAAGGAGA	TCTAACAACC	GACAACCATT	GGGTATGAAG												
	S	I	K	S	P	N	N	N	K	K	V	L	V	P	K	V	S
151	TCCATCAAGT	CTCCAAACAA	CAACAAGAAG	GTCTTGGTTC	CAAAGGTCTC												
	AGGTAGTTCA	GAGGTTTGT	GTTGTTCTTC	CAGAACCAAG	GTTTCCAGAG												
	G	L	Q	Y	R	V	F	R	V	R	L	P	D	P	N	K	F
201	TGGTTTGCAA	TACAGAGTCT	TCAGAGTCAG	ATTGCCAGAC	CCAAACAAGT												
	ACCAAACGTT	ATGTCTCAGA	AGTCTCAGTC	TAACGGTCTG	GGTTTGTTC												
	G	F	P	D	T	S	F	Y	N	P	D	T	Q	R	L	V	
251	TCGGTTTCCC	AGACACTTCC	TTCTACAACC	CAGACACTCA	AAGATTGGTC												
	AGCCAAAGGG	TCTGTGAAGG	AAGATGTTGG	GTCTGTGAGT	TTCTAACCAG												
	W	A	C	V	G	L	E	I	G	R	G	Q	P	L	G	V	G
301	TGGGCTTGTG	TCGGTTTGG	AATCGGTAGA	GGTCAACCAT	TGGGTGTTGG												
	ACCCGAACAC	AGCCAAACCT	TTAGCCATCT	CCAGTTGGTA	ACCCACAACC												
	V	S	G	H	P	Y	F	N	K	F	D	D	T	E	T	S	N
351	TGTCTCTGGT	CACCCATACT	TCAACAAGTT	CGACGACACC	GAAACCTCCA												
	ACAGAGACCA	GTGGGTATGA	AGTTGTTCAA	GCTGCTGTGG	CTTTGGAGGT												
	R	Y	P	A	Q	P	G	S	D	N	R	E	C	L	S	M	
401	ACAGATACCC	AGCTCAACCA	GGTTCTGACA	ACAGAGAATG	TTTGTCCATG												
	TGTCTATGGG	TCGAGTTGGT	CCAAGACTGT	TGTCTCTTAC	AAACAGGTAC												
	D	Y	K	Q	T	Q	L	C	L	I	G	C	K	P	P	T	G
451	GACTACAAGC	AAACCCAATT	GTGTTTGATC	GGTTGTAAGC	CACCAACTGG												
	CTGATGTTTCG	TTTGGGTAA	CACAACTAG	CCAACATTTCG	GTGGTTGACC												
	E	H	W	G	K	G	V	A	C	N	N	N	A	A	A	T	D
501	TGAACACTGG	GGTAAGGGTG	TTGCTTGTA	CAACAACGCT	GCTGCTACCG												
	ACTTGTGACC	CCATTCCCAC	AACGAACATT	GTTGTTGCGA	CGACGATGGC												
	C	P	P	L	E	L	F	N	S	I	I	E	D	G	D	M	
551	ACTGTCCACC	ATTGGAATTG	TTCAACTCCA	TCATCGAAGA	CGGTGACATG												
	TGACAGGTGG	TAACCTTAAC	AAGTTGAGGT	AGTAGCTTCT	GCCACTGTAC												
	V	D	T	G	F	G	C	M	D	F	G	T	L	Q	A	N	K

FIG.2A

5/10

601 GTCGACACTG GTTTCGGTTG TATGGACTTC GGTACCTTGC AAGCTAACAA
CAGCTGTGAC CAAAGCCAAC ATACCTGAAG CCATGGAACG TTCGATTGTT
S D V P I D I C N S T C K Y P D Y
651 GTCCGACGTT CCAATCGACA TCTGTAATC CACCTGTAAG TACCCAGACT
CAGGCTGCAA GGTTAGCTGT AGACATTGAG GTGGACATTC ATGGGTCTGA
L K M A S E P Y G D S L F F F L
701 ACTTGAAGAT GGCTTCTGAA CCATACGGTG ACTCCTTGTT CTTCTTCTTG
TGAACTTCTA CCGAAGACTT GGTATGCCAC TGAGGAACAA GAAGAAGAAC
R R E Q M F V R H F F N R A G K L
751 AGAAGAGAAC AAATGTTCTG CAGACACTTC TTCAACAGAG CTGGTAAGTT
TCTTCTCTTG TTTACAAGCA GTCTGTGAAG AAGTTGTCTC GACCATTCAA
G E A V P D D L Y I K G S G N T A
801 GGGTGAAGCT GTTCCAGACG ACTTGACAT CAAGGGTTCT GGTAACACCG
CCCACTTCGA CAAGGTCTGC TGAACATGTA GTTCCCAAGA CCATTGTGGC
V I Q S S A F F P T P S G S M V
851 CTGTCATCCA ATCCTCTGCT TTCTTCCCAA CTCCATCTGG TTCCATGGTC
GACAGTAGGT TAGGAGACGA AAGAAGGGTT GAGGTAGACC AAGGTACCAG
T S E S Q L F N K P Y W L Q R A Q
901 ACCTCTGAAT CTCAATTGTT CAACAAGCCA TACTGGTTGC AAAGAGCTCA
TGGAGACTTA GAGTTAACAA GTTGTTCCGT ATGACCAACG TTTCTCGAGT
G H N N G I C W G N Q L F V T V V
951 AGGTCACAAC AACGGTATCT GTTGGGGTAA CCAATTGTTC GTCACTGTCTG
TCCAGTGTTG TTGCCATAGA CAACCCCAT TGGTTAACAAG CAGTGACAGC
D T T R S T N M T L C T E V T K
1001 TCGACACCAC TAGATCCACT AACATGACCT TGTGTACCGA AGTCACCAAG
AGCTGTGGTG ATCTAGGTGA TTGTACTGGA ACACATGGCT TCAGTGGTTC
E G T Y K N D N F K E Y V R H V E
1051 GAAGGTACCT ACAAGAACGA CAACTTCAAG GAATACGTCA GACACGTCTGA
CTTCCATGGA TGTTCTTGCT GTTGAAGTTC CTTATGCAGT CTGTGCAGCT
E Y D L Q F V F Q L C K I T L T A
1101 GGAATACGAC TTGCAATTCG TCTTCCAATT GTGTAAGATC ACCTTGACTG
CCTTATGCTG AACGTTAAGC AGAAGGTTAA CACATTCTAG TGGAAGTAC
E I M T Y I H T M D S N I L E D
1151 CTGAAATCAT GACCTACATC CACACCATGG ACTCTAACAT CTTGGAAGAC
GACTTTAGTA CTGGATGTAG GTGTGGTACC TGAGATTGTA GAACCTTCTG
W Q F G L T P P P S A S L Q D T Y
1201 TGGCAATTCG GTTTGACTCC ACCACCATCT GCTTCCTTGC AAGACACCTA
ACCGTTAAGC CAACTGAGG TGGTGGTAGA CGAAGGAACG TTCTGTGGAT
R F V T S Q A I T C Q K T A P P K

FIG.2B


6/10

1251 CAGATTCGTC ACCTCTCAAG CTATCACCTG TCAAAAGACT GCTCCACCAA
GTCTAAGCAG TGGAGAGTTC GATAGTGGAC AGTTTTCTGA CGAGGTGGTT
E K E D P L N K Y T F W E V N L
1301 AGGAAAAGGA AGACCCATTG AACAAGTACA CCTTCTGGGA AGTCAACTTG
TCCTTTTCCT TCTGGGTAAC TTGTTTCATGT GGAAGACCCT TCAGTTGAAC
K E K F S A D L D Q F P L G R K F
1351 AAGGAAAAGT TCTCTGCTGA CTTGGACCAA TTCCCATTTG GTAGAAAGTT
TTCCTTTTCA AGAGACGACT GAACCTGGTT AAGGGTAACC CATCTTTCAA
L L Q S G L K A K P R L K R S A P
1401 CTTGTTGCAA TCTGGTTTGA AGGCTAAGCC AAGATTGAAG AGATCTGCTC
GAACAACGTT AGACCAAACCT TCCGATTTCG TTCTAACTTC TCTAGACGAG
T T R A P S T K R K K V K K * (SEQ ID NO:2)
1451 CAACCACTAG AGCTCCATCC ACCAAGAGAA AGAAGGTCAA GAAGTAA (SEQ ID NO:1)
GTTGGTGATC TCGAGGTAGG TGGTTCTCTT TCTTCCAGTT CTTTCATT (SEQ ID NO:10)

FIG.2C

7/10

Northern Blot of HPV 58 L1 wt and 58 L1 R transcripts.

**FIG.3**

Western Blot Analysis




FIG.4




FIG.5

10/10

Transmission EM of VLPs Composed of HPV 58 L1 R Protein Molecules.




FIG.6

SEQUENCE LISTING

<110> Merck & Co., Inc.

<120> OPTIMIZED EXPRESSION OF HPV 58 L1 In
YEAST

<130> 21561 PCT

<150> 60/519,211

<151> 2003-11-12

<160> 10

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1497

<212> DNA

<213> Artificial Sequence

<220>

<223> HPV58 L1-R

<400> 1

```

atgtccgtct ggagaccatc cgaagctacc gtctacttgc caccagttcc agtctccaag 60
gtcgtctcca ctgacgaata cgtctctaga acctctatct actactacgc tggttcctct 120
agattggttg ctggttggttaa cccatacttc tccatcaagt ctccaaacaa caacaagaag 180
gtcttggttc caaagggtctc tggtttgcaa tacagagtct tcagagtcag attgccagac 240
ccaaacaagt tcggtttccc agacacttcc ttctacaacc cagacactca aagattgggtc 300
tgggcttggt tcggtttgga aatcggtaga ggtcaaccat tgggtggttg tgtctctggt 360
cacccatact tcaacaagtt cgacgacacc gaaacctcca acagataccc agctcaacca 420
ggttctgaca acagagaatg tttgtccatg gactacaagc aaacccaatt gtgtttgatc 480
ggttgtaagc caccaactgg tgaacactgg ggtaagggtg ttgcttgtaa caacaacgct 540
gctgctaccg actgtccacc attggaattg ttcaactcca tcatcgaaga cggtgacatg 600
gtcgacactg gtttcggttg tatggacttc ggtaccttgc aagctaacaa gtccgacgtt 660
ccaatcgaca tctgtaactc cacctgtaag taccagact acttgaagat ggcttctgaa 720
ccatacggtg actccttggt cttcttcttg agaagagAAC aaatgttcgt cagacacttc 780
ttcaacagag ctggtaagtt ggggtgaagc gttccagacg acttgtaacat caagggttct 840
ggtaacaccg ctgtcatcca atcctctgct ttcttcccaa ctccatctgg ttccatgggtc 900
acctctgaat ctcaattggt caacaagcca tactgggtgc aaagagctca aggtcacaac 960
aacggtatct gttggggtaa ccaattgttc gtcactgtcg tcgacaccac tagatccact 1020
aacatgacct tgtgtaccga agtcaccaag gaaggtaact acaagaacga caacttcaag 1080
gaatacgtca gacacgtcga ggaatacgac ttgcaattcg tcttccaatt gtgtaagatc 1140
accttgactg ctgaaatcat gacctacatc cacaccatgg actctaacat cttggaagac 1200
tggaattcg gtttgactcc accaccatct gcttccttgc aagacaccta cagattcgtc 1260
acctctcaag ctatcacctg tcaaaagact gctccaccaa aggaaaagga agaccattg 1320
aacaagtaca ctttctggga agtcaacttg aaggaaaagt tctctgctga cttggaccaa 1380
ttcccatagg gtagaaagtt cttgttgcaa tctgggttga aggctaagcc aagattgaag 1440
agatctgctc caaccactag agctccatcc accaagagaa agaagggtcaa gaagtaa 1497

```

<210> 2

<211> 498

<212> PRT

<213> Human Papillomavirus Type 58

<400> 2

```

Met Ser Val Trp Arg Pro Ser Glu Ala Thr Val Tyr Leu Pro Pro Val
 1           5           10           15
Pro Val Ser Lys Val Val Ser Thr Asp Glu Tyr Val Ser Arg Thr Ser
          20          25          30

```

Ile	Tyr	Tyr	Tyr	Ala	Gly	Ser	Ser	Arg	Leu	Leu	Ala	Val	Gly	Asn	Pro
		35					40					45			
Tyr	Phe	Ser	Ile	Lys	Ser	Pro	Asn	Asn	Asn	Lys	Lys	Val	Leu	Val	Pro
	50					55					60				
Lys	Val	Ser	Gly	Leu	Gln	Tyr	Arg	Val	Phe	Arg	Val	Arg	Leu	Pro	Asp
65					70					75					80
Pro	Asn	Lys	Phe	Gly	Phe	Pro	Asp	Thr	Ser	Phe	Tyr	Asn	Pro	Asp	Thr
				85					90					95	
Gln	Arg	Leu	Val	Trp	Ala	Cys	Val	Gly	Leu	Glu	Ile	Gly	Arg	Gly	Gln
			100					105					110		
Pro	Leu	Gly	Val	Gly	Val	Ser	Gly	His	Pro	Tyr	Phe	Asn	Lys	Phe	Asp
		115					120					125			
Asp	Thr	Glu	Thr	Ser	Asn	Arg	Tyr	Pro	Ala	Gln	Pro	Gly	Ser	Asp	Asn
	130					135					140				
Arg	Glu	Cys	Leu	Ser	Met	Asp	Tyr	Lys	Gln	Thr	Gln	Leu	Cys	Leu	Ile
145					150					155					160
Gly	Cys	Lys	Pro	Pro	Thr	Gly	Glu	His	Trp	Gly	Lys	Gly	Val	Ala	Cys
				165					170					175	
Asn	Asn	Asn	Ala	Ala	Ala	Thr	Asp	Cys	Pro	Pro	Leu	Glu	Leu	Phe	Asn
			180					185					190		
Ser	Ile	Ile	Glu	Asp	Gly	Asp	Met	Val	Asp	Thr	Gly	Phe	Gly	Cys	Met
		195					200					205			
Asp	Phe	Gly	Thr	Leu	Gln	Ala	Asn	Lys	Ser	Asp	Val	Pro	Ile	Asp	Ile
	210					215					220				
Cys	Asn	Ser	Thr	Cys	Lys	Tyr	Pro	Asp	Tyr	Leu	Lys	Met	Ala	Ser	Glu
225					230					235					240
Pro	Tyr	Gly	Asp	Ser	Leu	Phe	Phe	Phe	Leu	Arg	Arg	Glu	Gln	Met	Phe
				245					250					255	
Val	Arg	His	Phe	Phe	Asn	Arg	Ala	Gly	Lys	Leu	Gly	Glu	Ala	Val	Pro
			260					265					270		
Asp	Asp	Leu	Tyr	Ile	Lys	Gly	Ser	Gly	Asn	Thr	Ala	Val	Ile	Gln	Ser
		275					280					285			
Ser	Ala	Phe	Phe	Pro	Thr	Pro	Ser	Gly	Ser	Met	Val	Thr	Ser	Glu	Ser
	290					295					300				
Gln	Leu	Phe	Asn	Lys	Pro	Tyr	Trp	Leu	Gln	Arg	Ala	Gln	Gly	His	Asn
305					310					315					320
Asn	Gly	Ile	Cys	Trp	Gly	Asn	Gln	Leu	Phe	Val	Thr	Val	Val	Asp	Thr
				325					330					335	
Thr	Arg	Ser	Thr	Asn	Met	Thr	Leu	Cys	Thr	Glu	Val	Thr	Lys	Glu	Gly
			340					345					350		
Thr	Tyr	Lys	Asn	Asp	Asn	Phe	Lys	Glu	Tyr	Val	Arg	His	Val	Glu	Glu
		355					360					365			
Tyr	Asp	Leu	Gln	Phe	Val	Phe	Gln	Leu	Cys	Lys	Ile	Thr	Leu	Thr	Ala
	370					375					380				
Glu	Ile	Met	Thr	Tyr	Ile	His	Thr	Met	Asp	Ser	Asn	Ile	Leu	Glu	Asp
385					390					395					400
Trp	Gln	Phe	Gly	Leu	Thr	Pro	Pro	Pro	Ser	Ala	Ser	Leu	Gln	Asp	Thr
				405					410					415	
Tyr	Arg	Phe	Val	Thr	Ser	Gln	Ala	Ile	Thr	Cys	Gln	Lys	Thr	Ala	Pro
			420					425					430		
Pro	Lys	Glu	Lys	Glu	Asp	Pro	Leu	Asn	Lys	Tyr	Thr	Phe	Trp	Glu	Val
		435					440					445			
Asn	Leu	Lys	Glu	Lys	Phe	Ser	Ala	Asp	Leu	Asp	Gln	Phe	Pro	Leu	Gly
		450				455					460				
Arg	Lys	Phe	Leu	Leu	Gln	Ser	Gly	Leu	Lys	Ala	Lys	Pro	Arg	Leu	Lys
465					470					475					480
Arg	Ser	Ala	Pro	Thr	Thr	Arg	Ala	Pro	Ser	Thr	Lys	Arg	Lys	Lys	Val
				485					490					495	
Lys	Lys														

<210> 3
 <211> 1497
 <212> DNA
 <213> Human Papillomavirus Type 58

<400> 3
 atgtccgtgt ggcggcctag tgaggccact gtgtacctgc ctctgtgtgc tgtgtctaag 60
 gttgtaagca ctgatgaata tgtgtcacgc acaagcattt attattatgc tggcagttcc 120
 agacttttgg ctggttgcaa tccatatttt tccatcaaaa gtcccaataa caataaaaaa 180
 gtattagttc ccaaggatc aggcttacag tatagggtct ttaggggtgcg ttacacctgat 240
 cccaataaat ttgggtttcc tgatacatct tttataacc ctgatacaca acgtttgggc 300
 tgggcatgtg taggccttga aataggtagg ggacagccat tgggtgttgg cgtaagtggc 360
 catccttatt tcaataaatt tgatgacact gaaaccagta acagatatcc cgcacagcca 420
 gggctctgata acagggaatg cttatctatg gattataaac aaacacaatt atgtttaatt 480
 ggctgtaaac ctcccactgg tgagcattgg ggtaaagggt ttgcctgtaa caataatgca 540
 gctgctactg attgtcctcc attggaactt tttaattcta ttattgagga tggtagacatg 600
 gtagatacag ggtttgatg catggacttt ggtacattgc aggctaataa aagtgatgtg 660
 cctattgata tttgtaacag tacatgcaaa tatccagatt atttaaaaat ggccagtga 720
 ccttatgggg atagtgtgt cttttttctt agacgtgagc agatgtttgt taggcacttt 780
 tttaataggg ccggaact tgggcagggt gtcccggatg acctttatat taaagggtcc 840
 ggtaatactg cagttatcca aagtagtgca ttttttccaa ctctagtgg ctctatgggt 900
 acctcagaat cacaattatt taataagcct tattggctac agcgtgcaca aggtcataac 960
 aatggcattt gctggggcaa tcagttattt gttaccgtag ttgataccac tcgtagcact 1020
 aatatgacat tatgactga agtaactaag gaaggtagat ataaaaatga taattttaag 1080
 gaatatgtac gtcattgtga agaataatgac ttacagtttg tttttcagct ttgcaaaatt 1140
 aactaactg cagagataat gacatatata catactatgg attccaatat tttggaggac 1200
 tggcaatttg gtttaacacc tcctccgtct gccagtttac aggacacata tagatttggt 1260
 acctcccagg ctattacttg ccaaaaaaca gcacccccta aagaaaagga agatccatta 1320
 aataaatata ctttttgga ggtaactta aaggaaaagt tttctgcaga tctagatcag 1380
 tttcctttgg gacgaaagt tttattacaa tcaggcctta aagcaaagcc cagactaaaa 1440
 cgttcggccc ctactaccg tgcaccaacc accaaacgca aaaaggttaa aaaataa 1497

<210> 4
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

<400> 4
 atgtccgtgt ggcggcctag t 21

<210> 5
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

<400> 5
 gagatctgtg taagtaccac aacaatta 28

<210> 6
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

<400> 6
gggatccccac aaaacaaaat gtccgtgtgg c 31

<210> 7
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 7
gggatccgtg taagtaccac aacaatta 28

<210> 8
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 8
ggatcccaca aaacaaaatg tctgtctgga gacc 34

<210> 9
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 9
ggatcccaca aaacaaaatg tctgtctgga gacc 34

<210> 10
<211> 1497
<212> DNA
<213> Artificial Sequence

<220>
<223> HPV 58 L1 antisense

<400> 10
tacaggcaga cctctggtag gcttcgatgg cagatgaacg gtgggtcaagg tcagagggttc 60
cagcagaggt gactgcttat gcagagatct tggagataga tgatgatgcg accaaggaga 120
tctaacaacc gacaaccatt gggatgaag aggtagttca gaggtttgtt gttgttcttc 180
cagaaccaag gtttccagag accaaacggt atgtctcaga agtctcagtc taacgggtctg 240
ggtttgttca agccaaaggg tctgtgaagg aagatgttgg gtctgtgagt ttctaaccag 300
acccgaacac agccaaacct ttagccatct ccagttggta acccacaacc acagagacca 360
gtgggtatga agttgttcaa gctgctgtgg ctttggagggt tgtctatggg tcgagttggt 420
ccaagactgt tgtctcttac aaacagggtac ctgatgttcg tttggggttaa cacaactag 480
ccaacattcg gtgggtgacc acttgtgacc ccattcccac aacgaacatt gttgttgca 540
cgacgatggc tgacaggtgg taaccttaac aagttgaggt agtagcttct gccactgtac 600
cagctgtgac caaagccaac atacctgaag ccatggaacg ttcgattggt caggctgcaa 660
ggttagctgt agacattgag gtggacattc atgggtctga tgaacttcta ccgaagactt 720
ggtatgccac tgaggaacaa gaagaagaac tcttctcttg tttacaagca gtctgtgaag 780
aagttgtctc gaccattcaa cccacttcga caaggtctgc tgaacatgta gttcccaaga 840
ccattgtggc gacagtaggt taggagacga aagaagggtt gaggtagacc aaggtaccag 900
tggagactta gagttaacaa gttgttcggt atgaccaacg tttctcgagt tccagtgttg 960
ttgccataga caacccatt ggtaacaag cagtgcacgc agctgtgggtg atctaggtga 1020

ttgtactgga	acacatggct	tcagtgggtc	cttccatgga	tggtccttgct	ggtgaagttc	1080
cttatgcagt	ctgtgcagct	ccttatgctg	aacgttaagc	agaagggttaa	cacattctag	1140
tggaactgac	gacttttagta	ctggatgtag	gtgtgggtacc	tgagattgta	gaaccttctg	1200
accgttaagc	caaactgagg	tggtggtaga	cgaaggaacg	ttctgtggat	gtctaagcag	1260
tgagagagttc	gatagtggac	agttttctga	cgagggtggtt	tccttttcct	tctgggtaac	1320
ttgttcatgt	ggaagaccct	tcagttgaac	ttccttttca	agagacgact	gaacctgggtt	1380
aagggttaacc	catctttcaa	gaacaacggt	agaccaaact	tccgattcgg	ttctaacttc	1440
tctagacgag	gttggtgatc	tcgaggtagg	tggttctctt	tcttccagtt	cttcatt	1497

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☒ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.